arXiv:1001.1196v2 [math.AC] 11 Mar 2010

Bivariate Lagrange Interpolation on Tower Interpolation Sites *

Tian Dong [a] Tao Chen [b] Shugong Zhang [a] Xiaoying Wang [a]

[a] *School of Mathemathics, Key Lab. of Symbolic Computation and Knowledge Engineering (Ministry of Education), Jilin University, Changchun 130012, China*

[b] *School of Statistics, Capital University of Economics and Business, Beijing 100070, China*

Abstract

As is well known, the geometry of the interpolation site of a multivariate polynomial interpolation problem constitutes a dominant factor for the structures of the interpolation polynomials. Solving interpolation problems on interpolation sites with special geometries in theory may be a key step to the development of general multivariate interpolation theory. In this paper, we introduce a new type of 2-dimensional interpolation sites, tower interpolation sites, whose associated degree reducing Lagrange interpolation monomial and Newton bases w.r.t. fixed standard term orders such as lexicographical order, total degree lexicographical order, etc. can be figured out theoretically. Inputting these interpolation bases into Buchberger-Möller(BM) algorithm, we can also construct the reduced Gröbner bases for related vanishing ideals. Experimental results show that in this way we can get the bases much faster than inputting tower sites directly into BM algorithm.

Key words: Bivariate Lagrange interpolation, Degree reducing interpolation space, Tower interpolation site, Gröbner basis

---

1. Introduction

Let $\mathbb{F}$ be an arbitrary field and $\mathbb{F}_q$ a finite prime field of size q. $\Pi^d := \mathbb{F}[x_1, x_2, \ldots, x_d]$ stands for the d-variate polynomial ring over $\mathbb{F}$. Given a set $\Xi = \{\xi^{(1)}, \ldots, \xi^{(\mu)}\} \subset \mathbb{F}^d$ of μ distinct points. For prescribed values $f_i \in \mathbb{F}, i = 1, \ldots, \mu$, find all polynomials $p \in \Pi^d$ such that

$$p(\xi^{(i)}) = f_i, \quad i = 1, \ldots, \mu. \tag{1}$$

---

* Dedicated to Professor Wen-Tsun Wu on the occasion of his ninetieth birthday.

**This research was partially supported by the National Grand Fundamental Research 973 Program of China(No. 2004CB318000) and the National Natural Science Foundation of China (Nos. 10601020).

Email addresses: `dongtian@jlu.edu.cn` (Tian Dong), `chentao@cueb.edu.cn` (Tao Chen), `sgzh@jlu.edu.cn` (Shugong Zhang), `wang.xiao.ying@163.com` (Xiaoying Wang).

Preprint submitted to Elsevier *12 March 2010*

The problem is called *d-variate Lagrange interpolation*, and the points are named *interpolation nodes*, Ξ the *interpolation site*, and p an *interpolation polynomial.* Generally, we only search interpolation polynomials in a subspace $\mathcal{P} \subset \Pi^d$. If there is a unique $p \in \mathcal{P}$ satisfying (1) for any fixed f_i's, we call $\mathcal{P}$ an *interpolation space* that gives rise to a *regular*(or *poised*) *interpolation scheme* $(\Xi, \mathcal{P})$(see Lorentz (1992)).

As a basic subject in Approximation Theory and Numerical Analysis, multivariate Lagrange interpolation has many applications in various fields of mathematics and computer science. Nevertheless, unlike univariate Lagrange interpolation, it is a relatively new topic with many basic problems unsolved. Since the set of all polynomials in Π^d vanishing at Ξ constitutes the vanishing ideal $\mathcal{I}(\Xi)$ of Ξ, multivariate Lagrange interpolation becomes a particular example of *ideal interpolation* introduced in Birkhoff (1979) and surveyed by de Boor (2005). Recent years, there has been considerable interest in the topic, for example, cf. Sauer and Xu (1995); Bos et al. (2007); Shekhtman (2009) etc.

In most cases, especially from a numerical point of view, we are not interested in all polynomials satisfying (1) but the one of "minimal degree", as in the univariate cases. In Sauer (2006), the notion of *degree reducing interpolation spaces* w.r.t. a fixed term order $\prec$ for interpolation problem (1) was introduced. Let $\mathcal{P}_\prec(\Xi) \subset \Pi^d$ be the subspace spanned by the Gröbner éscalier $\mathrm{N}_\prec(\mathcal{I}(\Xi))$ of $\mathcal{I}(\Xi)$ w.r.t. $\prec$. de Boor (2007) pointed out that $\mathcal{P}_\prec(\Xi)$ is a canonical $\prec$-degree reducing interpolation space as the unique one spanned by monomials. When $d = 1$, since $\mathcal{I}(\Xi) = \langle (x - \xi^{(1)})(x - \xi^{(2)}) \cdots (x - \xi^{(\mu)}) \rangle$, $\mathcal{P}_\prec(\Xi) = \mathrm{Span}_\mathbb{F}\{x^k : 0 \le k \le \mu - 1\}$ is a classical one. But when $d > 1$, we can not obtain $\mathcal{P}_\prec(\Xi)$ theoretically for any Ξ any more since the geometry of Ξ, as a dominant factor for $\mathrm{N}_\prec(\mathcal{I}(\Xi))$, becomes very complex. In these cases, $\mathcal{P}_\prec(\Xi)$ can only be acquired by Buchberger-Möller(BM for short) algorithm or variations thereof.

As a milestone algorithm for the computation of vanishing ideals, BM algorithm was presented in Möller and Buchberger (1982). For any interpolation site Ξ and fixed term order $\prec$, BM algorithm yields the reduced Gröbner basis for $\mathcal{I}(\Xi)$ w.r.t. $\prec$ and the monomial basis(i.e. $\mathrm{N}_\prec(\mathcal{I}(\Xi))$) as well as a Newton basis for $\mathcal{P}_\prec(\Xi)$. However, as we all known, BM algorithm(and the variations) has a poor complexity that limits its applications. One reason for this may be that the geometry of Ξ is not concerned by these algorithms. Therefore, if we can solve the interpolation problems on some special distributed nodes in theory, general interpolation theory will certainly benefit from it and the algorithms will also be improved for some cases.

In this decade, Sauer (2004); Crainic (2004); Chen et al. (2006) studied multivariate Lagrange interpolation on a special type of interpolation sites, lower interpolation sites, and constructed the associated degree reducing interpolation monomial and Newton bases in theory. Enlightened by them, this paper introduces a new type of bivariate interpolation sites, tower interpolation sites, whose geometry is more complex than lower sites nonetheless the related degree reducing interpolation bases can still be obtained theoretically.

In this paper, we will study bivariate Lagrange interpolation on tower interpolation sites. After introducing tower sites and comparing them with lower sites in Section 3, we will present the degree reducing interpolation bases w.r.t. several standard term orders in Section 4 as main results. The last section, Section 5, will give our improved BM algorithm for tower sites and illustrate some experimental results.

Next comes Section 2 that serves as a preparation for the paper.

2. Preliminaries

For the reader's convenience, this section will introduce some notation and recall some basic facts.

Let $\mathbb{N}_0$ stand for the monoid of nonnegative integers. A *polynomial* $f \in \Pi^2$ is of the form

$$f = \sum_{(i,j)\in\mathbb{N}_0^2} f_{ij} x^i y^j, \qquad \#\{(i,j) \in \mathbb{N}_0^2 : 0 \neq f_{ij} \in \mathbb{F}\} < \infty.$$

The set of bivariate monomials in Π^2 is denoted by $\mathbb{T}^2$.

Fix a term order $\prec$ on Π^2 that may be lexicographical order $\prec_{\mathrm{lex}}$, inverse lexicographical order $\prec_{\mathrm{inlex}}$, total degree lexicographical order $\prec_{\mathrm{tdlex}}$, or total degree inverse lexicographical order $\prec_{\mathrm{tdinlex}}$ etc. (see Becker and Weispfenning (1993)). For all nonzero $f \in \Pi^2$, we let $\mathrm{LT}(f)$ signify the *leading term*,$\mathrm{LM}(f)$ the *leading monomial*, and $\mathrm{LC}(f)$ the *leading coefficient* of f. Furthermore, for a non-empty subset $F \subset \Pi^2$, put

$$\mathrm{LT}(F) := \{\mathrm{LT}(f) : f \in F\}.$$

Let $\mathcal{A}$ be a finite subset of $\mathbb{N}_0^2$. $\mathcal{A}$ is called a *lower* set if, for any $(i,j) \in \mathcal{A}$, we always have

$$\{(i',j') \in \mathbb{N}_0^2 : 0 \leq i' \leq i, 0 \leq j' \leq j\} \subset \mathcal{A}. \tag{2}$$

Especially, $(0,0) \in \mathcal{A}$. Let $m_j = \max_{(h,j)\in\mathcal{A}} h, 0 \leq j \leq \nu = \max_{(0,k)\in\mathcal{A}} k$. Obviously, $\mathcal{A}$ can be determined uniquely by the ordered $(\nu+1)$-tuple $(m_0, m_1, \ldots, m_\nu)$ hence represented as $\mathrm{L}_x(m_0, m_1, \ldots, m_\nu)$. In like manner, we can also represent $\mathcal{A}$ as $\mathrm{L}_y(n_0, n_1, \ldots, n_{m_0})$ with $n_i = \max_{(i,k)\in\mathcal{A}} k, 0 \leq i \leq m_0$. It is easy to see that $\nu = n_0$.

Definition 1 (Crainic (2004))**.** We say that an interpolation site Ξ in $\mathbb{F}^2$ is *lower* if there exists a lower set $\mathcal{A} \subset \mathbb{N}_0^2$ such that Ξ can be written as

$$\Xi = \{(x_i, y_j) : (i,j) \in \mathcal{A}\}, \tag{3}$$

where the x_i's are distinct numbers, and similarly the y_j's. We also say that Ξ is $\mathcal{A}$-lower.

In de Boor (2007), the $\prec$*−degree* of a polynomial $f \in \Pi^2$ was defined to be the leading bidegree w.r.t. $\prec$

$$\delta(f) := (i,j), \quad x^i y^j = \mathrm{LM}(f),$$

with $\delta(0)$ undefined. For any $f, g \in \Pi^2$, if $\delta(f) \prec \delta(g)$ then we say that f is of *lower degree* than g and use the abbreviation

$$f \prec g := \delta(f) \prec \delta(g).$$

In addition, $f \preceq g$ is interpreted as the degree of f is lower than or equal to that of g.

Definition 2 (Sauer (2006))**.** Fix term order $\prec$. We call a subspace $\mathcal{P} \subset \Pi^2$ a *degree reducing interpolation space* w.r.t. $\prec$ for the bivariate Lagrange interpolation (1) if

(i) $\mathcal{P}$ is an *interpolation space*.

(ii) $\mathcal{P}$ is $\prec$*−reducing*, i.e., when $L_{\mathcal{P}}$ denotes the Lagrange projector with range $\mathcal{P}$, then the interpolation polynomial

$$L_{\mathcal{P}} q \preceq q, \qquad \forall q \in \Pi^2.$$

If an interpolation space $\mathcal{P}$ for (1) is degree reducing w.r.t. $\prec$, a basis $\{p_1, \dots, p_\mu\}$ for $\mathcal{P}$ will be called a *degree reducing interpolation basis* w.r.t. $\prec$ for (1). Assume that $p_1 \prec p_2 \prec \cdots \prec p_\mu$. If

$$p_j(\xi^{(i)}) = \delta_{ij}, \quad 1 \le i \le j \le \mu,$$

for some suitable reorder of Ξ, then we call the basis a *degree reducing interpolation Newton basis*(DRINB) w.r.t. $\prec$ for (1). Let $G_\prec$ be the reduced Gröbner basis for the vanishing ideal $\mathcal{I}(\Xi)$ w.r.t.$\prec$. The set

$$\mathrm{N}_\prec(\mathcal{I}(\Xi)) := \{x^i y^j \in \mathbb{T}^2 : \mathrm{LT}(g) \nmid x^i y^j, \forall g \in G_\prec\} \tag{4}$$

is the *Gröbner éscalier* of $\mathcal{I}(\Xi)$ w.r.t. $\prec$. Recall Section 1. As the monomial basis for $\mathcal{P}_\prec(\Xi)$, $\mathrm{N}_\prec(\mathcal{I}(\Xi))$ is named the *degree reducing interpolation monomial basis*(DRIMB) w.r.t. $\prec$ for (1).

Finally, we will redescribe BM algorithm with the notation established above.

Algorithm 1. (BM Algorithm)

Input: An interpolation site $\Xi = \{\xi^{(i)} : i = 1, \dots, \mu\} \subset \mathbb{F}^d$ and a fixed term order $\prec$.

Output: The 3-tuple (G, N, Q), where G is the reduced Gröbner basis for $\mathcal{I}(\Xi)$ w.r.t. $\prec$, N is the Gröbner éscalier of $\mathcal{I}(\Xi)$ (the DRIMB for (1) also) w.r.t. $\prec$, and Q is a DRINB w.r.t. $\prec$ for (1).

BM1. Start with lists $G = [\ ], N = [\ ], Q = [\ ], L = [1]$, and a matrix $B = (b_{ij})$ over $\mathbb{F}$ with μ columns and zero rows initially.

BM2. If $L = [\ ]$, return (G, N, Q) and stop. Otherwise, choose the monomial $t = \min_\prec L$, and delete t from L.

BM3. Compute the evaluation vector $(t(\xi^{(1)}), \dots, t(\xi^{(\mu)}))$, and reduce it against the rows of B to obtain

$$(v_1, \dots, v_\mu) = (t(\xi^{(1)}), \dots, t(\xi^{(\mu)})) - \sum_i a_i(b_{i1}, \dots, b_{i\mu}), \quad a_i \in \mathbb{F}.$$

BM4.. If $(v_1, \dots, v_\mu) = (0, \dots, 0)$, then append the polynomial $t - \sum_i a_i q_i$ to the list G, where q_i is the ith element of Q. Remove from L all the multiples of t. Continue with **BM2**.

BM5. Otherwise $(v_1, \dots, v_\mu) \ne (0, \dots, 0)$, add $(v_1, \dots, v_\mu)$ as a new row to B and $t - \sum_i a_i q_i$ as a new element to Q. Append the monomial t to N, and add to L those elements of $\{x_1 t, \dots, x_d t\}$ that are neither multiples of an element of L nor of $\mathrm{LT}(G)$. Continue with **BM2**.

3. Tower interpolation sites

Given an interpolation site $\Xi \subset \mathbb{F}^2$. In Crainic (2004), two particular lower sets are constructed from Ξ, denoted by $S_x(\Xi), S_y(\Xi)$, which reflect the geometry of Ξ in certain sense.

Concretely, we cover the nodes in Ξ by lines $l_0^x, l_1^x, \dots, l_\nu^x$ parallel to the x-axis and assume that, without loss of generality, there are $m_j + 1$ nodes, say $u_{0j}^x, u_{1j}^x, \dots, u_{m_j,j}^x$, on l_j^x with $m_0 \ge m_1 \ge \cdots \ge m_\nu \ge 0$ hence the ordinates of u_{ij}^x and $u_{i'j}^x, i \ne i'$, same. Now, we define

$$S_x(\Xi) := \{(i, j) : 0 \le i \le m_j,\ 0 \le j \le \nu\}, \tag{5}$$

which clearly equals to $\mathrm{L}_x(m_0, m_1, \ldots, m_\nu)$. Similarly, we can also cover the nodes by lines $l_0^y, l_1^y, \ldots, l_\lambda^y$ parallel to the y-axis and denote the nodes on line l_i^y by $u_{i0}^y, u_{i1}^y, \ldots, u_{i,n_i}^y$ with $n_0 \geq n_1 \geq \cdots \geq n_\lambda \geq 0$ hence the abscissae of u_{ij}^y and $u_{ij'}^y, j \neq j'$, same. Here, we put

$$S_y(\Xi) := \{(i,j) : 0 \leq i \leq \lambda,\ 0 \leq j \leq n_i\} = \mathrm{L}_y(n_0, n_1, \ldots, n_\lambda).$$

In addition, we can also define the sets of abscissae and ordinates

$$\begin{aligned} H_j(\Xi) :=& \{\bar{x} : (\bar{x}, \bar{y}) \in l_j^x \cap \Xi\}, \quad 0 \leq j \leq \nu, \\ V_i(\Xi) :=& \{\bar{y} : (\bar{x}, \bar{y}) \in l_i^y \cap \Xi\}, \quad 0 \leq i \leq \lambda. \end{aligned} \tag{6}$$

Now, we will introduce our key notion: tower interpolation site.

Definition 3. We say that an interpolation site Ξ in $\mathbb{F}^2$ is *x-tower* if lower set $S_x(\Xi) = \mathrm{L}_x(m_0, m_1, \ldots, m_\nu) \subset \mathbb{N}_0^2$ with $m_0 > m_1 > \cdots > m_\nu \geq 0$ such that

$$\Xi := \{(x_{ij}, y_j) : (i,j) \in S_x(\Xi)\}, \tag{7}$$

where $x_{ij} \in H_0(\Xi), (i,j) \in S_x(\Xi)$, are distinct for fixed j. We also call Ξ a $S_x(\Xi)$-*x-tower* interpolation site. In like fashion, if lower set $S_y(\Xi) = \mathrm{L}_y(n_0, n_1, \ldots, n_\lambda) \subset \mathbb{N}_0^2$, $n_0 > n_1 > \cdots > n_\lambda \geq 0$, such that

$$\Xi := \{(x_i, y_{ij}) : (i,j) \in S_y(\Xi)\}, \tag{8}$$

where $y_{ij} \in V_0(\Xi), (i,j) \in S_y(\Xi)$, are distinct for fixed i, we will call Ξ a *y-tower* interpolation site.

We have mentioned in Section 1, Definition 3 is enlightened by the notion of lower sites. Next, we will compare them in detail. Recall Definition 1. We have the following criterion.

Theorem 4 (Crainic (2004)). *An interpolation site $\Xi \subset \mathbb{F}^2$ is lower if and only if $S_x(\Xi) = S_y(\Xi)$.*

Let Ξ be an $\mathcal{A}$-lower site. By Definition 1 and Theorem 4, we can deduce that $\mathcal{A} = S_x(\Xi) = S_y(\Xi)$. Hence, (3) can be redescribed as

$$\Xi := \{(x_i, y_j) : (i,j) \in S_x(\Xi)\} \tag{9}$$

or

$$\Xi := \{(x_i, y_j) : (i,j) \in S_y(\Xi)\}. \tag{10}$$

Observing (7), (9) and (8), (10), we find that they are similar in form each to each nevertheless it seems that tower sites make the nodes on a line much freer than lower sites, which is supported by the following lemma as an alternative criterion for lower sites.

Lemma 5. *Resume the notation established above. An interpolation site $\Xi \subset \mathbb{F}^2$ is lower if and only if*

$$H_0(\Xi) \supseteq H_1(\Xi) \supseteq \cdots \supseteq H_\nu(\Xi) \tag{11}$$

or

$$V_0(\Xi) \supseteq V_1(\Xi) \supseteq \cdots \supseteq V_\lambda(\Xi). \tag{12}$$

Proof. Assume that interpolation site Ξ is lower. By (5), (6), and (9), we have $H_j(\Xi) = \{x_0, x_1, \ldots, x_{m_j}\}, j = 0, \ldots, \nu$. Since $m_0 \geq m_1 \geq \cdots \geq m_\nu \geq 0$, (11) follows. (12) can be proved in like manner.

Conversely, we assume that (11) holds. Suppose $S_x(\Xi) = \mathrm{L}_x(m_0, m_1, \ldots, m_\nu) = \mathrm{L}_y(n'_0, n'_1, \ldots, n'_{\lambda'})$ and $S_y(\Xi) = \mathrm{L}_y(n_0, n_1, \ldots, n_\lambda)$. Since $S_x(\Xi)$ is lower, we have $\lambda' = m_0$ and $n'_0 = \nu$. By (6), $\#H_j(\Xi) = m_j + 1, j = 0, \ldots, \nu$.

Next, we shall verify that $\lambda = m_0 = \lambda'$, namely we can cover Ξ by exactly $m_0 + 1$ vertical lines $l^y_i, i = 0, \ldots, m_0$, which pass the $m_0 + 1$ nodes on line l^x_0 respectively. Prove this by contradiction. Since $\lambda \geq m_0$ apparently, equality can fail only when $\lambda > m_0$, i.e., there exists at least one node $(\bar{x}, \bar{y}) \in \Xi$ such that $(\bar{x}, \bar{y}) \notin l^y_i, i = 0, \ldots, m_0$. Because $(\bar{x}, \bar{y}) \in \Xi$, from (6), there must exist some $H_{j^*}(\Xi), 0 \leq j^* \leq \nu$, such that $\bar{x} \in H_{j^*}(\Xi)$. (11) implies that $\bar{x} \in H_0(\Xi)$, which contradicts $(\bar{x}, \bar{y}) \notin l^y_i, i = 0, \ldots, m_0$.

Finally, we will prove

$$n'_h = n_h, \quad h = 0, \ldots, m_0, \tag{13}$$

which implies $S_x(\Xi) = S_y(\Xi)$ immediately. We prove it by induction on h. When $h = 0$, for any node $(x^{(\nu)}, y^{(\nu)}) \in l^x_\nu$, $x^{(\nu)} \in H_\nu(\Xi)$ hence $x^{(\nu)} \in H_j(\Xi), 0 \leq j \leq \nu$, due to (11). So we have found $\nu+1$ nodes on line $x = x^{(\nu)}$ that obviously equals to some $l^y_{i^*}, 0 \leq i^* \leq m_0$. From the definition of ν and $S_y(\Xi)$, we have $n_0 = n'_0 = \nu$, namely (13) is true for $h = 0$. Now, we assume (13) for $0 \leq h \leq k < m_0$. When $h = k + 1$, by the induction hypothesis, we can find $k + 1$ lines, without loss of generality, $l^y_0, \ldots, l^y_k$ that contain exactly $n_h + 1$ nodes, $0 \leq h \leq k$, of Ξ respectively. Since $S_x(\Xi) = \mathrm{L}_y(n'_0, n'_1, \ldots, n'_{m_0})$, we have $(k + 1, n'_{k+1}) \in S_x(\Xi)$ which implies that there must exist at least one node $(x^{(n'_{k+1})}, y^{(n'_{k+1})}) \in l^x_{n'_{k+1}}$ which is not on $l^y_0, \ldots, l^y_k$. By (11), we can find $n'_{k+1} + 1$ nodes on line $x = x^{(n'_{k+1})}$ that is a member of $\{l^y_{k+1}, \ldots, l^y_{m_0}\}$ hence $n_{k+1} \geq n'_{k+1}$. From the definition of $S_y(\Xi)$, there exists one line in $\{l^y_{k+1}, \ldots, l^y_{m_0}\}$ that have exactly $n_{k+1} + 1$ nodes distributed among some lines of $\{l^x_0, \ldots, l^x_{n_{k+1}}\}$ due to (11). Therefore, we have $n'_{k+1} \geq n_{k+1}$ that leads to $n_{k+1} = n'_{k+1}$, namely (13) holds for $h = k + 1$. Consequently, $S_x(\Xi) = \mathrm{L}_y(n'_0, n'_1, \ldots, n'_{m_0}) = \mathrm{L}_y(n_0, n_1, \ldots, n_{m_0}) = S_y(\Xi)$ which implies that Ξ is lower due to Theorem 4.

Swapping the roles of x and y, we can also prove the other statement similarly therefore complete the proof. $\square$

Note that it is easy for Lemma 5 to extend to higher dimensions but difficult for Theorem 4.

Now, we go back to tower sites. For an x-tower site Ξ, although $H_j(\Xi) \subsetneq H_0(\Xi), j = 1, \ldots, \nu$, it may have nothing to do with any other $H_i(\Xi), i = 1, \ldots, j-1, j+1, \ldots, \nu$. Comparing with (11), we find that, on one line, lower sites allow free amounts of nodes while tower sites allow free positions of nodes. The following proposition reveals the relation between them.

Proposition 6. *Resume the notation established above. If Ξ satisfies $H_0(\Xi) \supsetneq H_1(\Xi) \supsetneq \cdots \supsetneq H_\nu(\Xi)$, then Ξ is lower if and only if it is x-tower. Similarly, if Ξ satisfies $V_0(\Xi) \supsetneq V_1(\Xi) \supsetneq \cdots \supsetneq V_\lambda(\Xi)$, then Ξ is lower iff it is y-tower.*

Proof. Assume that Ξ is lower with $S_x(\Xi) = \mathrm{L}_x(m_0, m_1, \ldots, m_\nu)$. Since $H_0(\Xi) \supsetneq H_1(\Xi) \supsetneq \cdots \supsetneq H_\nu(\Xi)$, by (6), we have $m_0 > m_1 > \cdots > m_\nu \geq 0$ which implies that

Ξ is a $S_x(\Xi)$-x-tower site according to (7). Conversely, if Ξ is x-tower, since $H_0(\Xi) \supsetneq H_1(\Xi) \supsetneq \cdots \supsetneq H_\nu(\Xi)$, by Lemma 5, Ξ is lower follows.

In very like fashion, we can prove the other statement. □

Proposition 6 shows that the notions of lower sites and tower sites are not mutually exclusive as there exist lower sites that are tower sites, but there are also tower sites are not lower and vice versa. The following examples show them with illustrations.

Example 7. Observe (a), (b), (c) of Fig. 1. Ξ_1 is an x-tower site that is not lower while Ξ_2 is a lower site that is not x-tower or y-tower. Ξ_3 is a y-tower site that is lower too. (d) of Fig. 1 illustrates $S_y(\Xi_3)$ that is equal to $S_x(\Xi_3)$.



(a) Ξ_1

(b) Ξ_2

(c) Ξ_3

(d) $S_y(\Xi_3)$

Fig. 1. Illustrations for Example 7

4. Main Results

In this section, we will pursue the degree reducing interpolation bases w.r.t. a fixed term order for bivariate Lagrange interpolation on a tower site Ξ. Since an interpolation polynomial of minimal total degree is prefered in most situations, the $\prec_{\text{tdlex}}$ and $\prec_{\text{tdinlex}}$ cases will be discussed at the beginning. Note that we will continue with the notation established in the previous sections.

Lemma 8 (Dong et al. (2005))**.** *Let* $\Xi = \{(x_0, y_0), (x_1, y_0), \ldots, (x_m, y_0)\} \subset \mathbb{F}^2$ *be a set of distinct nodes on line* $y = y_0$*. Then*

$$\mathcal{I}(\Xi) = \langle (x - x_0)(x - x_1) \cdots (x - x_m), y - y_0 \rangle.$$

As a main result of the paper, the following theorem gives the DRIMB w.r.t. $\prec_{\text{tdlex}}$ or $\prec_{\text{tdinlex}}$ for bivariate Lagrange interpolation on an x-tower or y-tower site respectively in theory.

Theorem 9. *Given an* x*-tower site* $\Xi \subset \mathbb{F}^2$*. The DRIMB w.r.t.* $\prec_{\text{tdlex}}$ *for* (1) *is*

$$N_x = \{x^i y^j : (i, j) \in S_x(\Xi)\}.$$

If Ξ *is* y*-tower, then the DRIMB w.r.t.* $\prec_{\text{tdinlex}}$ *for* (1) *is*

$$N_y = \{x^i y^j : (i, j) \in S_y(\Xi)\}.$$

Proof. We will only prove the first statement. The second one can be verified in very like fashion.

For fixed j, by Lemma 8, the ideal $\mathcal{I}_j := \mathcal{I}(\{(x_{0j}, y_j), (x_{1j}, y_j), \ldots, (x_{m_j,j}, y_j)\})$, $j = 0, \ldots, \nu$, equals

$$\langle (x - x_{0j})(x - x_{1j}) \cdots (x - x_{m_j,j}), y - y_j \rangle.$$

Obviously, $\mathcal{I}_j$'s are pairwise comaximal hence the ideal $\mathcal{I}(\Xi)$ is equal to

$$\bigcap_{j=0}^{\nu} \mathcal{I}_j = \prod_{j=0}^{\nu} \mathcal{I}_j = \prod_{j=0}^{\nu} \langle (x - x_{0j})(x - x_{1j}) \cdots (x - x_{m_j,j}), y - y_j \rangle.$$

Let G be the reduced Gröbner basis for $\mathcal{I}(\Xi)$ and G_k the reduced Gröbner basis for $\prod_{j=0}^{k} \mathcal{I}_j$ w.r.t. $\prec_{\text{tdlex}}$, $0 \le k \le \nu$. We will use induction on ν to prove

$$\mathrm{LT}(G) = \{x^{m_0+1}, x^{m_1+1}y, x^{m_2+1}y^2, \ldots, x^{m_\nu+1}y^\nu, y^{\nu+1}\}. \tag{14}$$

First of all, since

$$\mathcal{I}_0 = \langle (x - x_{0,0})(x - x_{1,0}) \cdots (x - x_{m_0,0}), y - y_0 \rangle,$$

(14) is true for $\nu = 0$.

Now, let $\nu = 1$. Ξ is x-tower implies that $m_0 > m_1$ and $H_1(\Xi) \subsetneq H_0(\Xi)$. Therefore,

$$\begin{aligned}\mathcal{I}_0 \cdot \mathcal{I}_1 =& \langle (x - x_{0,0}) \cdots (x - x_{m_0,0}), y - y_0 \rangle \cdot \langle (x - x_{0,1}) \cdots (x - x_{m_1,1}), y - y_1 \rangle \\ =& \langle (x - x_{0,0}) \cdots (x - x_{m_0,0}), (x - x_{0,0}) \cdots (x - x_{m_0,0})(y - y_1), \\ & (x - x_{0,1}) \cdots (x - x_{m_1,1})(y - y_0), (y - y_0)(y - y_1) \rangle \\ =& \langle (x - x_{0,0}) \cdots (x - x_{m_0,0}), (x - x_{0,1}) \cdots (x - x_{m_1,1})(y - y_0), (y - y_0)(y - y_1) \rangle.\end{aligned}$$

It is easy to check that $\mathrm{LT}(G_1) = \{x^{m_0+1}, x^{m_1+1}y, y^2\}$ which means that (14) holds for $\nu = 1$.

When $\nu = 2$, since $m_1 > m_2$ and $H_2(\Xi) \subsetneq H_0(\Xi)$, we get, after some easy computations,

$$\begin{aligned}\mathcal{I}_0 \cdot \mathcal{I}_1 \cdot \mathcal{I}_2 =&\langle (x - x_{0,0}) \cdots (x - x_{m_0,0}), (y - y_0)(y - y_1)(y - y_2),\\ &(x - x_{0,1}) \cdots (x - x_{m_1,1})(x - x_{0,2}) \cdots (x - x_{m_2,2})(y - y_0),\\ &(x - x_{0,1}) \cdots (x - x_{m_1,1})(y - y_0)(y - y_2),\\ &(x - x_{0,2}) \cdots (x - x_{m_2,2})(y - y_0)(y - y_1)\rangle\\ =:&\langle g_0^{(2)}, g_1^{(2)}, g_2^{(2)}, g_3^{(2)}, g_4^{(2)}\rangle.\end{aligned}$$

Noticing that $m_1 > m_2$, we let $\hat{q}, \hat{r} \in \Pi^1$ be the quotient and remainder respectively of the division of $(x - x_{0,1}) \cdots (x - x_{m_1,1})$ by $(x - x_{0,2}) \cdots (x - x_{m_2,2})$, namely

$$(x - x_{0,1}) \cdots (x - x_{m_1,1}) = \hat{q}(x - x_{0,2}) \cdots (x - x_{m_2,2}) + \hat{r}.$$

Denote the remainder of $g_3^{(2)}$ w.r.t. $g_4^{(2)}$ by $\bar{g}_3^{(2)}$. One can check readily that

$$\bar{g}_3^{(2)} = g_3^{(2)} - \hat{q}g_4^{(2)}.$$

On the other hand, since

$$g_2^{(2)} = \frac{1}{y_1 - y_2}\left[(x - x_{0,2}) \cdots (x - x_{m_2,2})g_3^{(2)} - (x - x_{0,1}) \cdots (x - x_{m_1,1})g_4^{(2)}\right],$$

we have $g_2^{(2)} \xrightarrow{\{g_3^{(2)}, g_4^{(2)}\}}_{+} 0$. It follows that

$$\mathcal{I}_0 \cdot \mathcal{I}_1 \cdot \mathcal{I}_2 =\langle\ g_0^{(2)}, g_1^{(2)}, \bar{g}_3^{(2)}, g_4^{(2)}\rangle.$$

We claim that

$$G_2 = G_2' := \{g_0^{(2)}, g_1^{(2)}, (y_1 - y_2)^{-1}\bar{g}_3^{(2)}, g_4^{(2)}\},$$

where $y_1 - y_2 = \mathrm{LC}(\bar{g}_3^{(2)})$, i.e., $(y_1 - y_2)^{-1}\bar{g}_3^{(2)}$ is monic.

In fact, if $S(f, g)$ stands for the S-polynomial of arbitrary polynomials $f, g \in \Pi^2$, $S(g_0^{(2)}, g_1^{(2)}) \xrightarrow{G_2'}_{+} 0$ follows immediately because $\mathrm{LM}(g_0^{(2)})$ and $\mathrm{LM}(g_1^{(2)})$ are relatively prime. Notice that $(x - x_{0,2}) \cdots (x - x_{m_2,2})$ is a factor of $(x - x_{0,0}) \cdots (x - x_{m_0,0})$, we get

$$\begin{aligned}S(g_0^{(2)}, g_4^{(2)}) =&y^2 g_0^{(2)} - x^{m_0-m_2}g_4^{(2)}\\ =&y^2 g_0^{(2)} - x^{m_0-m_2}g_4^{(2)} - (y - y_0)(y - y_1)g_0^{(2)} + (y - y_0)(y - y_1)g_0^{(2)}\\ =&y^2 g_0^{(2)} - x^{m_0-m_2}g_4^{(2)} - (y - y_0)(y - y_1)g_0^{(2)} + \frac{(x - x_{0,0}) \cdots (x - x_{m_0,0})}{(x - x_{0,2}) \cdots (x - x_{m_2,2})}g_4^{(2)}\\ =&((y_0 + y_1)y - y_0y_1)g_0^{(2)} + \left(\frac{(x - x_{0,0}) \cdots (x - x_{m_0,0})}{(x - x_{0,2}) \cdots (x - x_{m_2,2})} - x^{m_0-m_2}\right)g_4^{(2)}.\end{aligned}$$

It is easy to check that $\mathrm{LM}(S(g_0^{(2)}, g_4^{(2)})) = x^{m_0+1}y$. Since

$$\mathrm{LM}(((y_0 + y_1)y - y_0y_1)g_0^{(2)}) = x^{m_0+1}y,$$

$$\mathrm{LM}\left(\left(\frac{(x - x_{0,0}) \cdots (x - x_{m_0,0})}{(x - x_{0,2}) \cdots (x - x_{m_2,2})} - x^{m_0-m_2}\right)g_4^{(2)}\right) = x^{m_0}y^2,$$

we have

$$\begin{aligned}\mathrm{LM}(S(g_0^{(2)},g_4^{(2)})) = \max_{\prec_{\mathrm{tdlex}}}\Bigg(&\mathrm{LM}(((y_0+y_1)y-y_0y_1)g_0^{(2)}),\\ &\mathrm{LM}\left(\left(\frac{(x-x_{0,0})\cdots(x-x_{m_0,0})}{(x-x_{0,2})\cdots(x-x_{m_2,2})}-x^{m_0-m_2}\right)g_4^{(2)}\right)\Bigg)\\ =&x^{m_0+1}y,\end{aligned}$$

which implies that $S(g_0^{(2)},g_4^{(2)}) \xrightarrow{G_2'}_+ 0$. Similarly, by

$$S(g_1^{(2)},g_4^{(2)}) = (x^{m_2+1}-(x-x_{0,2})\cdots(x-x_{m_2,2}))g_1^{(2)} - y_2g_4^{(2)},$$

and $\mathrm{LM}(S(g_1^{(2)},g_4^{(2)})) = x^{m_2+1}y^2$,

$$\begin{aligned}&\max_{\prec_{\mathrm{tdlex}}}\left(\mathrm{LM}((x^{m_2+1}-(x-x_{0,2})\cdots(x-x_{m_2,2}))g_1^{(2)}),\mathrm{LM}(-y_2g_4^{(2)})\right)\\ =&\max_{\prec_{\mathrm{tdlex}}}(x^{m_2}y^3,x^{m_2+1}y^2) = x^{m_2+1}y^2 = \mathrm{LM}(S(g_1^{(2)},g_4^{(2)}))\end{aligned}$$

follows, namely $S(g_1^{(2)},g_4^{(2)}) \xrightarrow{G_2'}_+ 0$.

In like manner, we can also prove that

$$S(g_0^{(2)},(y_1-y_2)^{-1}\bar{g}_3^{(2)}) \xrightarrow{G_2'}_+ 0, \quad S(g_1^{(2)},(y_1-y_2)^{-1}\bar{g}_3^{(2)}) \xrightarrow{G_2'}_+ 0.$$

Hence, there only remains $S((y_1-y_2)^{-1}\bar{g}_3^{(2)},g_4^{(2)}) \xrightarrow{G_2'}_+ 0$ to be checked. Actually, it can be deduced that $S((y_1-y_2)^{-1}\bar{g}_3^{(2)},g_4^{(2)})$ equals

$$\frac{\hat{r}}{y_1-y_2}g_1^{(2)} + y_1\cdot(y_1-y_2)^{-1}\bar{g}_3^{(2)} + \left(\hat{q}-x^{m_1-m_2}\right)g_4^{(2)}.$$

Observing the structure of $S((y_1-y_2)^{-1}\bar{g}_3^{(2)},g_4^{(2)})$, we know that its leading monomial is $x^{m_1+1}y$, therefore, from

$$\begin{aligned}\mathrm{LM}\left(\frac{\hat{r}}{y_1-y_2}g_1^{(2)}\right) &= x^{m_2}y^3,\\ \mathrm{LM}(y_1\cdot(y_1-y_2)^{-1}\bar{g}_3^{(2)}) &= x^{m_1+1}y,\\ \mathrm{LM}\left(\left(\hat{q}-x^{m_1-m_2}\right)g_4^{(2)}\right) &= x^{m_1}y^2,\end{aligned}$$

$S((y_1-y_2)^{-1}\bar{g}_3^{(2)},g_4^{(2)}) \xrightarrow{G_2'}_+ 0$.

The above arguments lead to the conclusion: S-polynomial $S(g,g') \xrightarrow{G_2'}_+ 0$ for all $g,g' \in G_2', g \neq g'$. Consequently, from Buchberger's S-pair criterion, G_2' is a Gröbner basis for $\mathcal{I}_0\cdot\mathcal{I}_1\cdot\mathcal{I}_2$ w.r.t. $\prec_{\mathrm{tdlex}}$.

Moreover, for any polynomial $g \in G_2'$, it is evident that

(1) $\mathrm{LC}(g) = 1$,

(2) No monomial of g lies in $\langle \mathrm{LT}(G_2'-\{g\})\rangle$,

which implies that G_2' is the reduced Gröbner basis for $\mathcal{I}_0\cdot\mathcal{I}_1\cdot\mathcal{I}_2$ w.r.t. $\prec_{\mathrm{tdlex}}$. Hence (14) holds for $\nu = 2$.

Now, for $\nu = k$, assume (14). That is to say

$$\mathrm{LT}(G_k) = \{x^{m_0+1}, x^{m_1+1}y, x^{m_2+1}y^2, \dots, x^{m_k+1}y^k, y^{k+1}\}.$$

Let $G_k = \{g_0^{(k)}, \dots, g_{k+1}^{(k)}\}$. Without loss of generality, we may assume that

$$\begin{aligned} \mathrm{LT}(g_i^{(k)}) =& x^{m_i+1}y^i, \quad i = 0, \dots, k, \\ \mathrm{LT}(g_{k+1}^{(k)}) =& y^{k+1}, \end{aligned}$$

which imply that

$$\begin{aligned} g_0^{(k)} &= (x - x_{0,0}) \cdots (x - x_{m_0,0}), \\ g_{k+1}^{(k)} &= (y - y_0)(y - y_1) \cdots (y - y_k). \end{aligned}$$

When $\nu = k+1$, since Ξ is x-tower, we obtain

$$\begin{aligned} \prod_{i=0}^{k+1} \mathcal{I}_i =& (\prod_{i=0}^{k} \mathcal{I}_i)\mathcal{I}_{k+1} \\ =& \langle\ g_0^{(k)}, \dots, g_{k+1}^{(k)} \rangle \cdot \langle (x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}), y - y_{k+1} \rangle \\ =& \langle\ g_0^{(k+1)}, g_0^{(k)}(y - y_{k+1}), \\ & \quad g_1^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}), g_1^{(k)}(y - y_{k+1}), \\ & \quad \cdots \\ & \quad g_{k-1}^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}), g_{k-1}^{(k)}(y - y_{k+1}), \\ & \quad g_k^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}), g_k^{(k)}(y - y_{k+1}), \\ & \quad g_{k+1}^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}), g_{k+1}^{(k)}(y - y_{k+1}) \rangle, \end{aligned}$$

where $g_0^{(k+1)} = g_0^{(k)}$. Thus, $g_0^{(k)}(y - y_{k+1})$ can be removed from the ideal basis above. By the induction hypothesis, we have

$$g_{k+2}^{(k+1)} := g_{k+1}^{(k)}(y - y_{k+1}) = (y - y_0)(y - y_1) \cdots (y - y_{k+1}).$$

We denote polynomial $g_{k+1}^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1})$ by $g_{k+1}^{(k+1)}$. For $E_1 := \{g_{k+1}^{(k+1)}\}$, suppose $g_k^{(k)}(y - y_{k+1}) \xrightarrow{E_1}_+ g_k^{(k+1)}$. Since $m_k > m_{k+1}$, we have

$$\mathrm{LM}(g_k^{(k+1)}) = x^{m_k+1}y^k.$$

Furthermore, recalling the case $\nu = 2$, we can check easily that

$$g_k^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}) \xrightarrow{F_1}_+ 0,$$

where $F_1 := \{g_{k+1}^{(k+1)}, g_k^{(k+1)}\}$, which implies that the polynomial can be removed from the basis.

Let $E_2 := F_1$ and suppose $g_{k-1}^{(k)}(y - y_{k+1}) \xrightarrow{E_2}_+ g_{k-1}^{(k+1)}$. We can also deduce that $\mathrm{LM}(g_{k-1}^{(k+1)}) = x^{m_{k-1}+1}y^{k-1}$ and

$$g_{k-1}^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}) \xrightarrow{F_2}_+ 0,$$

where $F_2 := \{g_{k+1}^{(k+1)}, g_{k-1}^{(k+1)}\}$.

In this way we can construct two sequences $E_1, E_2, \ldots, E_k$ and $F_1, F_2, \ldots, F_k$ such that

$$g_i^{(k)}(y - y_{k+1}) \xrightarrow{E_{k+1-i}}_+ g_i^{(k+1)},$$
$$g_i^{(k)}(x - x_{0,k+1}) \cdots (x - x_{m_{k+1},k+1}) \xrightarrow{F_{k+1-i}}_+ 0,$$

where, for $i = 1, \ldots, k$,

$$E_i = \{g_{k+2-i}^{(k+1)}, \ldots, g_{k+1}^{(k+1)}\},$$
$$F_i = \{g_{k+1}^{(k+1)}, g_{k+1-i}^{(k+1)}\}.$$

Let $\bar{g}_i^{(k+1)} = \mathrm{LC}(g_i^{(k+1)})^{-1} g_i^{(k+1)}$. With quite similar methods that we used in the case $\nu = 2$, we can prove finally that

$$\{\bar{g}_0^{(k+1)}, \bar{g}_1^{(k+1)}, \ldots, \bar{g}_{k+1}^{(k+1)}\}$$

is the reduced Gröbner basis for $\prod_{i=0}^{k+1} \mathcal{I}_i$ w.r.t. $\prec_{\text{tdlex}}$, i.e., (14) holds when $\nu = k + 1$. By (4) and the definition of $S_x(\Xi)$, we have

$$\mathrm{N}_{\prec_{\text{tdlex}}}(\mathcal{I}(\Xi)) = N_x = \{x^i y^j : (i, j) \in S_x(\Xi)\},$$

which complete the proof. □

The next two theorems present degree reducing interpolation Newton bases w.r.t. $\prec_{\text{tdlex}}$ or $\prec_{\text{tdinlex}}$ for bivariate Lagrange interpolation on an x-tower or y-tower site respectively.

Theorem 10. *Let $\Xi \subset \mathbb{F}^2$ be an x-tower site of nodes $u_{mn}^x = (x_{mn}, y_n), (m, n) \in S_x(\Xi)$, which gives rise to polynomials*

$$\phi_{ij}^x = \varphi_{ij}^x \prod_{t=0}^{j-1} (y - y_t) \prod_{s=0}^{i-1} (x - x_{sj}), \quad (i, j) \in S_x(\Xi),$$

where $\varphi_{ij}^x = 1 / \prod_{t=0}^{j-1} (y_j - y_t) \prod_{s=0}^{i-1} (x_{ij} - x_{sj}) \in \mathbb{F}$, and the empty products are taken as 1. Then

$$Q_x = \{\phi_{ij}^x : (i, j) \in S_x(\Xi)\} \tag{15}$$

is a DRINB *w.r.t. $\prec_{\text{tdlex}}$ for* (1) *satisfying*

$$\phi_{ij}^x(u_{mn}^x) = \delta_{(i,j),(m,n)}, \quad (i, j) \succeq_{\text{inlex}} (m, n).$$

Proof. Fix $(i, j) \in S_x(\Xi)$. If $(i, j) = (m, n)$, by $y_0 \neq y_1 \neq \cdots \neq y_j$ and $x_{0j} \neq x_{1j} \neq \cdots \neq x_{ij}$, we have

$$\phi_{ij}^x(u_{ij}^x) = \varphi_{ij}^x \prod_{t=0}^{j-1} (y_j - y_t) \prod_{s=0}^{i-1} (x_{ij} - x_{sj}) = \varphi_{ij}^x / \varphi_{ij}^x = 1.$$

Otherwise, if $(i, j) \succ_{\text{inlex}} (m, n)$, we have $j > n$, or $j = n, i > m$. When $j > n$, we have

$$\phi_{ij}^x(u_{mn}^x) = \varphi_{ij}^x (y_n - y_0) \cdots (y_n - y_n) \cdots (y_n - y_{j-1}) \prod_{s=0}^{i-1} (x_{mn} - x_{sj}) = 0,$$

and when $j = n, i > m$,

$$\begin{aligned}\phi_{ij}^x(u_{mn}^x) &= \varphi_{ij}^x \prod_{t=0}^{j-1}(y_n - y_t)(x_{mn} - x_{0j})\cdots(x_{mn} - x_{mj})\cdots(x_{mn} - x_{i-1,j})\\ &= \varphi_{ij}^x \prod_{t=0}^{n-1}(y_n - y_t)(x_{mn} - x_{0n})\cdots(x_{mn} - x_{mn})\cdots(x_{mn} - x_{i-1,n})\\ &= 0,\end{aligned}$$

which leads to

$$\phi_{ij}^x(u_{mn}^x) = 0, \quad (i,j) \succ_{\mathrm{inlex}} (m,n),$$

namely Q_x is a Newton basis for $\mathrm{Span}_{\mathbb{F}} Q_x$. By Theorem 9, it is easy to check that $\mathrm{Span}_{\mathbb{F}} Q_x = \mathrm{Span}_{\mathbb{F}} N_x = \mathcal{P}_{\prec_{\mathrm{tdlex}}}(\Xi)$. Therefore, Q_x is a DRINB w.r.t. $\prec_{\mathrm{tdlex}}$ for (1). $\square$

Similarly, we can prove the following theorem:

Theorem 11. *Let $\Xi \subset \mathbb{F}^2$ be a y-tower site of nodes $u_{mn}^y = (x_m, y_{mn}), (m,n) \in S_y(\Xi)$. We define the polynomials*

$$\phi_{ij}^y = \varphi_{ij}^y \prod_{s=0}^{i-1}(x - x_s) \prod_{t=0}^{j-1}(y - y_{it}), \quad (i,j) \in S_y(\Xi),$$

where $\varphi_{ij}^y = 1/\prod_{s=0}^{i-1}(x_i - x_s)\prod_{t=0}^{j-1}(y_{ij} - y_{it}) \in \mathbb{F}$. The empty products are taken as 1. Then,

$$Q_y = \{\phi_{ij}^x : (i,j) \in S_y(\Xi)\} \tag{16}$$

is a DRINB *w.r.t. $\prec_{\mathrm{tdinlex}}$ for* (1) *satisfying*

$$\phi_{ij}^y(u_{mn}^y) = \delta_{(i,j),(m,n)}, \quad (i,j) \succeq_{\mathrm{lex}} (m,n).$$

Now, we will turn to $\prec_{\mathrm{lex}}$ and $\prec_{\mathrm{inlex}}$ cases. For any interpolation site $\Xi \subset \mathbb{F}^2$, the following theorem presents the degree reducing interpolation monomial and Newton bases w.r.t. $\prec_{\mathrm{lex}}$ or $\prec_{\mathrm{inlex}}$ for (1) theoretically.

Theorem 12 (Wang et al. (2009))**.** *Let*

$$\Xi = \{u_{mn}^x = (x_{mn}^x, y_{mn}^x) : (m,n) \in S_x(\Xi)\} = \{u_{mn}^y = (x_{mn}^y, y_{mn}^y) : (m,n) \in S_y(\Xi)\}$$

be an interpolation site in $\mathbb{F}^2$. Then

(i) *the set $N_x := \{x^i y^j : (i,j) \in S_x(\Xi)\}$ is the* DRIMB *as well as $Q_x := \{\phi_{ij}^x : (i,j) \in S_x(\Xi)\}$ is a* DRINB *w.r.t. $\prec_{\mathrm{lex}}$ for* (1)*, where*

$$\phi_{ij}^x = \varphi_{ij}^x \prod_{t=0}^{j-1}(y - y_{0t}^x) \prod_{s=0}^{i-1}(x - x_{sj}^x), \quad (i,j) \in S_x(\Xi),$$

with $\varphi_{ij}^x = 1/\prod_{t=0}^{j-1}(y_{0j}^x - y_{0t}^x)\prod_{s=0}^{i-1}(x_{ij}^x - x_{sj}^x) \in \mathbb{F}$ and the empty products taken as 1;

(ii) *the set $N_y := \{x^i y^j : (i,j) \in S_y(\Xi)\}$ is the* DRIMB *as well as $Q_y := \{\phi_{ij}^y : (i,j) \in S_y(\Xi)\}$ is a* DRINB *w.r.t. $\prec_{\mathrm{inlex}}$ for* (1)*, where*

$$\phi_{ij}^y = \varphi_{ij}^y \prod_{s=0}^{i-1}(x - x_{s0}^y) \prod_{t=0}^{j-1}(y - y_{it}^y), \quad (i,j) \in S_y(\Xi),$$

with $\varphi^y_{ij} = 1/\prod_{s=0}^{i-1}(x^y_{i0} - x^y_{s0})\prod_{t=0}^{j-1}(y^y_{ij} - y^y_{it}) \in \mathbb{F}$*, and the empty products are taken as 1.*

Since Theorem 12 holds for any interpolation site in $\mathbb{F}^2$, it obviously holds for tower sites. It is easy to find that the N_x in the theorem is same to the one in Theorem 9, namely the DRIMB w.r.t. $\prec_{\text{lex}}$ is same to the one w.r.t. $\prec_{\text{tdlex}}$ for interpolation on an x-tower site. In addition, the Q_x here is same to (15) when Ξ is x-tower. For N_y and Q_y, similar statements are also true for a y-tower site.

In a word, for a preassigned x-tower site, we can obtain the DRIMB w.r.t. $\prec_{\text{tdlex}}$ or $\prec_{\text{lex}}$ by Theorem 9, and a DRINB w.r.t. $\prec_{\text{tdlex}}$ or $\prec_{\text{lex}}$ by Theorem 10. The y-tower cases can also be solved by Theorem 9 and 11.

5. Algorithm and Experimental results

Let Ξ be a tower site in $\mathbb{F}^2$. Suppose (G, N, Q) be the 3-tuple output of BM algorithm. Theorem 9-11 present us N and Q theoretically hence we can obtain them with little effort. According to Marinari et al. (1993), $\text{LT}(G)$ is contained in the border set of N. Therefore, inputting N, Q into BM algorithm can make us obtain G much faster than compute G directly with BM algorithm.

Algorithm 2. (TBM algorithm)

Input: A x-tower(y-tower) interpolation site $\Xi \subset \mathbb{F}^2$ of μ nodes and fixed term order $\prec_{\text{tdlex}}(\prec_{\text{tdinlex}})$ or $\prec_{\text{lex}}(\prec_{\text{inlex}})$.

Output: The 3-tuple (G, N, Q), where G is the reduced Gröbner basis for $\mathcal{I}(\Xi)$, N is the Gröbner éscalier $\text{N}(\mathcal{I}(\Xi))$, and Q is a DRINB for (1).

TBM1. Construct lower set $S_x(\Xi)(S_y(\Xi))$ following the process introduced in Section 3.

TBM2. Compute the sets N and Q according to Theorem 9-11.

TBM3. Compute the border set $L := \{x \cdot t : t \in N\} \bigcup \{y \cdot t : t \in N\} \setminus N$.

TBM4. Construct $\mu \times \mu$ matrix B whose (h, k) entry is $\phi^x_h(u^x_k)(\phi^y_h(u^y_k))$ where $\phi^x_h(\phi^y_h)$, $u^x_k(u^y_k)$ are hth and kth elements of $Q = \{\phi^x_{ij}(\phi^y_{ij}) : (i,j) \in S_x(\Xi)(S_y(\Xi))\}$ and $\Xi = \{u^x_{mn}(u^y_{mn}) : (m,n) \in S_x(\Xi)(S_y(\Xi))\}$ w.r.t. the increasing $\prec_{\text{inlex}}(\prec_{\text{lex}})$ on (i,j) and (m,n) respectively.

TBM5. Goto **BM2** with N, Q, L, B for the reduced Gröbner basis G.

In the following, we will show the timings for the computations of BM-problems on tower sites in finite prime fields w.r.t. term order $\prec_{\text{inlex}}$ or $\prec_{\text{tdlex}}$. BM and TBM algorithms were implemented on Maple 12 installed on a laptop with 2 Gb RAM and 1.8 GHz CPU.

For field $\mathbb{F}_{23}$ and $\prec_{\text{tdlex}}$,

$\#\Xi$ Algorithms	100	300	500	900	1300
TBM	0.125 s	1.297 s	4.953 s	20.609 s	56.906 s
BM	0.531 s	9.437 s	53.625 s	224.812 s	861.531 s

For field $\mathbb{F}_{43}$ and $\prec_{\text{inlex}}$,

Algorithms \ #Ξ	600	900	1200	1500	1800
TBM	7.781 s	21.094 s	45.282 s	87.734 s	134.265 s
BM	121.094 s	362.000 s	1002.578 s	2077.907	2729.750 s

References

Becker, T., Weispfenning, V., 1993. Gröbner Bases. Vol. 141 of Graduate Texts in Mathematics. Springer-Verlag, New York.

Birkhoff, G., 1979. The algebra of multivariate interpolation. In: Coffman, C. V., Fix, G. J. (Eds.), Constructive Approaches to Mathematical Models. Academic Press, New York, pp. 345–363.

Bos, L., De Marchi, S., Vianello, M., Xu, Y., 2007. Bivariate Lagrange interpolation at the Padua points: The ideal theory approach. Numer. Math. 108 (1), 43–57.

Chen, T., Dong, T., Zhang, S., 2006. The Newton interpolation bases on lower sets. J. Inf. Comput. Sci. 3 (3), 385–394.

Crainic, N., 2004. Multivariate Birkhoff-Lagrange interpolation schemes and cartesian sets of nodes. Acta Math. Univ. Comenian.(N.S.) LXXIII (2), 217–221.

de Boor, C., 2005. Ideal interpolation. In: Chui, C. K., Neamtu, M., Schumaker, L. L. (Eds.), Approximation Theory XI: Gatlinburg 2004. Nashboro Press, Brentwood TN, pp. 59–91.

de Boor, C., 2007. Interpolation from spaces spanned by monomials. Adv. Comput. Math. 26 (1), 63–70.

Dong, T., Zhang, S., Lei, N., 2005. Interpolation basis for nonuniform rectangular grid. J. Inf. Comput. Sci. 2 (4), 671–680.

Lorentz, R., 1992. Multivariate Birkhoff Interpolation. Vol. 1516 of Lecture Notes in Mathematics. Springer, Heidelberg.

Marinari, M. G., Möller, H. M., Mora, T., 1993. Gröbner bases of ideals defined by functionals with an application to ideals of projective points. Appl. Algebra Engrg. Comm. Comput. 4 (2), 103–145.

Möller, H., Buchberger, B., 1982. The construction of multivariate polynomials with preassigned zeros. In: Calmet, J. (Ed.), Computer Algebra: EUROCAM '82. Vol. 144 of Lecture Notes in Computer Science. Springer, Berlin, pp. 24–31.

Sauer, T., 2004. Lagrange interpolation on subgrids of tensor product grids. Math. Comp. 73 (245), 181–190.

Sauer, T., 2006. Polynomial interpolation in several variables: Lattices, differences, and ideals. In: Jetter, K., Buhmann, M., Haussmann, W., Schaback, R., Stöckler, J. (Eds.), Topics in Multivariate Approximation and Interpolation. Vol. 12 of Studies in Computational Mathematics. Elsevier, Amsterdam, pp. 191–230.

Sauer, T., Xu, Y., 1995. On multivariate Lagrange interpolation. Math. Comp. 64 (211), 1147–1170.

Shekhtman, B., 2009. On the limits of Lagrange projectors. Constr. Approx. 29 (3), 293–301.

Wang, X., Zhang, S., Dong, T., 2009. A bivariate preprocessing paradigm for Buchberger-Möller algorithm, summited for Publication.